



Review

Ankylosing Spondylitis Associated Gene Polymorphism

Shuo Li ^{1,2,*}, Yu-Ming Wang ^{3,4,*}

¹Department of Laboratory Medicine, The Second Clinical Medical College, Yangtze University, Jingzhou, Hubei 434023, China; ²School of Basic Medicine, Health Science Center, Yangtze University, Jingzhou, Hubei 434023, China;

³Department of Spinal and Neural Function Reconstruction, Beijing Bo'ai Hospital, China Rehabilitation Research Center, Beijing 100068, China; ⁴Capital Medical University School of Rehabilitation Medicine, Beijing 100068, China.

* These authors contributed equally.

✉ Correspondence

Shuo Li, Department of Laboratory Medicine, the Second Clinical Medical College, Yangtze University, Jingzhou, Hubei 434023, China. Email: 1448927469@qq.com. Telephone number: 86-18372637673.

Received: May 5, 2020. **Accepted:** July 30, 2020. **Published online:** August 8, 2020. 10.46633

Cite this paper: Li, S. and Wang, Y.M. (2020) Ankylosing Spondylitis Associated Gene Polymorphism. *Global Journal of Immunology*, 1(1): 2-8. <http://naturescholars.com/gji.010102>. <https://doi.org/10.46633/gji.010102>.

Copyright © 2020 by Scholars Publishing, LLC.

Abstract

Ankylosing Spondylitis (AS) is an autoimmune arthritis disease, mostly found in 20-30 years old adolescents. Genetic factors play an important role in the pathogenesis of AS. It is generally believed that the disease is a polygenic hereditary disease and has strong correlation with human leukocyte antigen B27 (HLA-B27). But the mechanism is still unclear. In recent years, it has been found that endoplasmic reticulum aminopeptidase1 (ERAP-1) gene, tumor necrosis factor- α (TNF- α) gene, interleukin gene, and cytotoxic T lymphocyte associated antigen 4 (CTLA-4) gene are also correlated with the pathogenesis of AS. The research on these related genes can provide a basis for the diagnosis of AS. This article reviews the mechanism and polymorphism of these genes in AS.

Key words: Ankylosing spondylitis (AS); Human leukocyte antigen B27 gene (HLA-B27); Endoplasmic reticulum aminopeptidase1 gene (ERAP-1); Interleukin 23 receptor gene (IL-23R); Cytotoxic T lymphocyte associated antigen-4 gene (CTLA-4).

Introduction

AS is an autoimmune disease and chronic inflammatory disease with an estimated incidence

of 0.2-0.54% in Chinese Han population (1). It belongs to the category of rheumatoid diseases and is a kind of seronegative spondyloarthropathy. AS usually starts secretly and is easily neglected in the

early stage of the disease. Some AS patient's main features are the involvement of the spine and sacroiliac joints, who may have peripheral arthritis or peripheral joint adhesion (2). The typical manifestation of AS is inflammatory backache with varying degrees of vascular lesions. It has inflammation of tendons, ligaments and articular capsule fibers attached to bones, as well as extra articular manifestations such as colitis and psoriasis (3). The etiology of AS is unclear, but it may be caused by environmental factors based on genetic factors. It is generally believed that it is closely related to the expression of *HLA-B27*, but not all *HLA-B27* positive individuals develop into AS (4). In recent years, many other susceptibility genes have been found related to AS.

1. Human leukocyte antigen B27 gene (*HLA-B27*)

Human leukocyte antigen B27 (*HLA-B27*), encoded by the B gene, is located on chromosome 6 and belongs to I type MHC gene. Although *HLA-B27* plays an important role in the pathogenesis of AS, recently research have shown that *HLA-B27* accounts for only 20.1% of the total genetic susceptibility, and generally only 5% of *HLA-B27* positive individuals develop into AS in the Chinese Han population in Beijing(5). The pathogenesis of *HLA-B27* is still unclear, but there are mainly three hypotheses: the first theory is articular peptide: *HLA-B27* on APC binds to self-peptide or environmental polypeptide and then activates CD8+T cells; the second unconfirmed theory is the *HLA-B27* misfolding theory: the misfolding of *HLA-B27* in the endoplasmic reticulum results in endoplasmic reticulum stress and activates unfolded protein response, leading to the up-regulate of IL-23 in dendritic cells, which leads to inflammation; the third one is the cell surface homodimer hypothesis: *HLA-B27* folds abnormally on the surface of APC

to form homodimer, which can bind to killer immunoglobulin-like receptor 3DL2 (*KIR3DL2*)(6, 7). The mechanism is shown in step 1 and 2 of Figure 1. *HLA-B27* has many subtypes, some of which play an important role in AS. *HLA-B27:09* is not correlated with the incidence of AS (8). *HLA-B27:04*, *HLA-B27:05* and *HLA-B27:02* have the strongest correlation with AS and *HLA-B27:04* is the main subtype of AS in Chinese Han population in Beijing (5). Through genotyping of AS patients in Xinjiang, it is found that *HLA-B27:05* is the main subtype of Xinjiang Uygur, while *HLA-B27:04* is the main subtype of Xinjiang Han population (9). Different living habits, geographical environments and ethnic groups may produce different dominant subtypes of gene that affect the disease. By studying the subtypes of *HLA-B27* gene, we can diagnose the disease as soon as possible.

2. Endoplasmic reticulum aminopeptidase1 gene (*ERAP-1*)

ERAP-1 is another gene found mostly associated with AS after the discovery of *HLA-B27*. *ERAP-1* is an aminopeptidase located in the endoplasmic reticulum, and *ERAP-1* gene is located at chromosome 5q15 (5). *ERAP-1* is involved in the presentation processing of antigenic peptides. After the antigen peptides are pruned to the appropriate length for antigen presentation, they are bound to MHC-I molecules and migrate the cell membrane to induce an immune response. *ERAP-1* polymorphism changes the function of the enzyme and leads to the peptide expression and free heavy chain formation of *HLA-B27* (10). Studies have shown that there is interaction between *HLA-B27* gene and *ERAP-1* gene mutation, and that *ERAP-1* gene mutation is only associated with *HLA-B27* (+). *ERAP-1* abnormalities produce abnormal antigenic peptide spectrum, resulting in aberrant expression of *HLA-B27*. The mechanism is shown in step 3 of Figure 1. *rs27434* and *rs275825* polymorphisms of

ERAP-1 in HLA-B27:02 and HLA-B27:04 positive patients are associated with AS in Beijing area, but the *rs27434* is weakly associated with AS (5). According to Su's finding, *rs30183*, *rs27434*, *rs27044* have no correlation with AS in the Chinese Han population (11). However, *rs27044* polymorphism is associated with AS in a higher incidence of AS in CG genotype carriers by Tang's finding in Chinese Han population (12). This indicates that geographical factors may be an important reason for the difference. It is necessary to explore the correlation between *ERAP-1* gene polymorphism and the pathogenesis of AS through a lot of research and data analysis to provide more reasonable methods for the clinical diagnosis and treatment.

3. Interleukin 23 receptor gene (*IL-23R*)

Interleukin-23 receptor (*IL-23R*) is located on chromosome 1p31.3(13). Interleukin-23 receptor (*IL-23R*) encoded by *IL-23R* gene is a specific subunit of IL-23R/IL12R β 1 complex, that mediates the signal transduction of IL-23(14). IL-23R/IL-17 axis plays an important role in the pathogenesis of AS. IL-23, produced by dendritic cells and monocytes, promotes the polarization of activated T cells to helper Th17 cells and induces the production of IL-17A. IL-17A is pro-inflammatory cytokine, leading to bone destruction in AS patients (15). The mechanism is shown in step 4 of Figure 1. Changes in *IL-23R* gene polymorphism may cause dysfunction of the IL-23R/IL-17 axis, leading to the pathogenesis of AS. Many *IL-23R* gene polymorphisms have been discovered through a lot of research. By analyzing the polymorphism of *IL-23R rs10889677*, it is found that A allele is significantly correlated with the incidence of AS in European (13). So, *IL-23R rs10889677* polymorphism in the Asian remains to be further researched. *rs1004819*, *rs1495965* and *rs2201841* are susceptible to AS.

rs10489629, *rs1209026*, *rs11465804* and *rs1343151* have protective effects on AS, but *rs1209629* and *rs11465804* do not appear in Asian patients with AS (14). The different results may be due to the few numbers of cases and the small sample size, resulting in the limitation of experimental results. So, the undiscovered polymorphisms can be found through deeply research of *IL-23R* gene to provide a better method for the diagnosis of disease.

4. Cytotoxic T lymphocyte associated antigen-4 gene (*CTLA-4*)

CTLA-4 is also known as CD152. CTLA-4 is a transmembrane receptor on T cells and plays an important role in inhibiting the activation and tolerance of T cells. The *CTLA-4* gene is located in the 2q33 region of chromosome (16). T cells activation requires the co-stimulation of CD28 and CTLA-4. CD28 activates T cells, while CTLA-4 inhibits T cells activation by competing with CD28 for B7 in Algeria population (17). The mechanism is shown in step 5 of Figure 1. CTLA-4 regulates T cells activation through the balance of CD28 activation and CTLA-4 inhibition. After genotyping of *CTLA-4*, three high risk gene polymorphisms (*CTLA-4-318C/T*, *CTLA-4-CT60* and *CTLA-4+49A/G*) are found related to the pathogenesis of AS. The soluble CTLA-4 (sCTLA-4) in peripheral blood of patients with AS and normal subjects is detected. It is found that the sCTLA-4 in peripheral blood of AS patients is higher than normal controls, but the level of sCTLA-4 is not related to the severity of AS (18). *CTLA-4+49A/G* study shows that it has no significant correlation with the incidence of AS. It suggested that *CTLA-4+49A/G* might not be the main susceptibility gene in AS patients (19). Polymorphisms of *CTLA-4 +49A/G* and *CTLA-4-318T/C* are also not significantly correlated with the incidence of AS in Chen's finding(16). Genotyping of *CTLA-4/CT60* shows that the *CT60*G* allele is significantly correlated in women over 30

years old and the risk of AS is increased in HLA-B27 negative patients(17). In order to provide a new method for the diagnosis and treatment of AS in the future, the polymorphism of the *CTLA-4* gene needs further research.

5. Tumor necrosis factor- α gene (*TNF- α*)

Tumor necrosis factor- α (*TNF- α*) belongs to the TNF family, which is encoded by the *TNF- α* gene located on the chromosome 6p21.3 (20). TNF is secreted by macrophages and regulated by two distinct receptors, TNFR1 and TNFR2. TNFR1 promotes inflammation, while TNFR2 promotes neuroprotection in Russian Caucasian population (21). The mechanism is shown in step 6 of Figure 1. TNF is a key pro-inflammatory

cytokine in chronic inflammation and highly expressed in sacroiliac joint cells of AS patients in Chinese Han population (22). *TNF- α -238 GA/AA* and *TNF- α -308 GA/AA* genotype are found to be associated with improved spinal cord function (23). *TNF- α -1211c (rs1799964)* and *TNF- α -308 (rs1800629)* reduce the risk of AS in Chinese Han population, but *TNF- α -238 (rs361525)* is not associated with AS (22). The study of *TNF- α -308 (rs1800629)* gene in Norwegian population shows that it is also not associated with AS, but a new AS risk site is found in *TNF- α -238 (rs361525)* (24). The difference may be due to geographical factors, ethnic factors or too few samples. With further study of *TNF- α* , *TNF- α* inhibitor is a good drug for the treatment of AS in recent years.




Figure 1. Molecular Mechanism of Ankylosing Spondylitis

1. HLA-B27 on APC binds to peptide and activates CD8+T cells.
2. HLA-B27 folds abnormally on the surface of APC to form homologous dimer and binds to the killer immunoglobulin-like receptor 3DL2 (KIR3DL2).
3. ERAP-1 is involved in the processing of antigen peptide presentation. Abnormal ERAP-1 produces abnormal antigen peptide spectrum, resulting in abnormal expression of HLA-B27.
4. IL-23 promotes the polarization of activated T cells to the helper Th17 cells and induces the production of IL-17A by T helper 17 cells.
5. T cell activation requires the co-stimulation of CD28 and CTLA-4. CD28 activates T cells, while CTLA-4 inhibits T cell activation by competing with CD28 for B7.
6. TNF- α is secreted by macrophages and has pro-inflammatory effects.

Table1: Ankylosing spondylitis associated gene polymorphism

| Gene | Polymorphism | Mechanism | Result | References |
|---------------|--|---|---|------------|
| HLA-B27 | HLA-B27:05 B27:09 B27:04 B27:02 | HLA-B27 on APC binds to peptide and activates CD8+T cells. HLA-B27 folds abnormally on the surface of APC to form homologous dimer that binds to KIR3DL2. | HLA-B27:09 is not correlation with the incidence of AS. HLA-B27:04, HLA-B27:05 and HLA-B27:02 has the strongest correlation with AS in Beijing population. | 5-9 |
| ERAP-1 | rs27434 rs275825 rs30183 rs27044 | ERAP-1 takes part in the presentation of antigen peptide, pruning the peptide to the appropriate length of antigen presentation and then transferring to the cell membrane to activate the immune response. | rs27434 and rs275825 polymorphisms of ERAP-1 are associated with AS. rs30183 and rs27044 have no correlation with AS in Chinese Han population. | 5, 10-12 |
| IL-23R | rs1004819 rs1495965 rs2201841 rs10489629 | IL-23 produced by monocytes and dendritic cells, promotes the polarization of activates T cells to helper Th17 cells and induces IL-17A. | rs1004819, rs1495965 and rs2201841 are susceptible to AS. rs10489629 gene is decreased in AS. | 13-15 |
| CTLA-4 | -318*C/T +49*A/G CT60 | T cell activation requires the co-stimulation of CD28 and CTLA-4. CD28 activates T cells, while CTLA-4 inhibits T cell activation. | There is no correlation between CTLA-4+49A/G, CTLA-4-318T/C polymorphisms and the AS. The CT60*G allele increases the risk of AS in HLA-B27 negative group. | 16-19 |
| TNF- α | TNF- α -308 TNF- α -238 TNF- α -1211c | TNF is secreted by macrophages and regulated by TNFR1 and TNFR2. | TNF- α -1211c and TNF- α -308 reduce the risk of AS, but TNF- α -238 is not associated with AS. | 20-24 |

6. Other genes

In addition to the above genes, there are many other susceptible genes associated with the development of AS. *IL-10* gene is located in 1q31, and its *819C/T* polymorphism is significantly correlated with AS, but *592 C/A* polymorphism is not correlated with AS (25). By analyzing the polymorphism of *PON1* gene *rs662* and *rs854560*, it is found that the GG genotype and G allele of *rs662* polymorphism are closely related to the increased risk of AS, but the *rs854560* polymorphism is not associated with the onset of AS in Chinese Han population (26). The polymorphism of *PON1 rs662* gene can inhibit the activity of *PON1* and lead to the pathogenesis of AS. AS is also associated with Toll like receptor 4 (*TLR-4*) gene, transforming growth factor β 1 (*TGF- β 1*) and cytochrome P450 2D6 (*CYP2D6*). However, at present, the number of gene studies is

small, and needs to do a lot of research to verify it.

Conclusion

As a genetic associated disease, AS is closely related to many genes, such as *HLA-B27*, *ERAP-1*, *TNF- α* , *IL-23R* and *CTLA-4*. This review summarizes the mechanisms of these genes and draws the pictures above. In addition to genetic factors, AS may also be related to immunity, environment, drugs and infection and so on. We also need more and further studies of the polymorphism of AS susceptibility gene to understand the pathogenesis of AS, which can provide many new ways for the diagnosis and treatment of AS. It can also be used in accordance with the characteristics of gene polymorphism to make the clinical treatment individualized. It is also expected to find new susceptibility genes through many researches to provide more beneficial methods for clinical

diagnosis and treatment of AS.

Declarations

1) *Consent to publication*

We declare that all authors agreed to publish the manuscript at this journal

based on the signed Copyright Transfer Agreement and followed publication ethics.

2) *Ethical approval and consent to participants*

Not applicable.

3) *Disclosure of conflict of interests*

We declare that no conflict of interest exists.

4) *Funding*

None

5) *Availability of data and material*

We declare that the data supporting the results reported in the article are available in the published article.

6) *Authors' Contributions*

Authors contributed to this paper with the design (SL), literature search (SL), drafting (SL), revision (SL and YMW), editing (SL and YMW) and final approval (SL).

7) *Acknowledgement*

None

8) *Authors' biography*

None

References

- Li L, Shi BL, Zheng WK, Xing WH, Zhao Y, Li F, et al. Association of IL-1A and IL-1B polymorphisms with ankylosing spondylitis among the Chinese Han population: a case-control study. *Oncotarget*.2017;8(17):28278-84.
- Raychaudhuri SP, Deodhar A. The classification and diagnostic criteria of ankylosing spondylitis. *J Autoimmun*. 2014;48-49:128-33.
- Bergman M, Lundholm A. Managing morbidity and treatment-related toxicity in patients with ankylosing spondylitis. *Rheumatology (Oxford)*. 2018;57(3):419-28.
- V R, E G, MA B, RD I, N H. Pathogenesis of ankylosing spondylitis - recent advances and future directions. *D 101500080*. (1759-4804 (Electronic)): 359-67.
- Zhang Z, Dai D, Yu K, Yuan F, Jin J, Ding L, et al. Association of HLA-B27 and ERAP1 with ankylosing spondylitis susceptibility in Beijing Han Chinese. *Tissue Antigens*. 2014;83(5):324-9.
- Simone D, Al Mossawi MH, Bowness P. Progress in our understanding of the pathogenesis of ankylosing spondylitis. *Rheumatology (Oxford)*. 2018;57 (suppl_6):vi4-vi9.
- Bowness P. Hla-B27. *Annu Rev Immunol*.2015;33:29-48.
- Chen B, Li J, He C, Li D, Tong W, Zou Y, et al. Role of HLA-B27 in the pathogenesis of ankylosing spondylitis (Review). *Mol Med Rep*. 2017;15(4):1943-51.
- Schittenhelm RB, Sivaneswaran S, Lim Kam Sian TC, Croft NP, Purcell AW. Human Leukocyte Antigen (HLA) B27 Allotype-Specific Binding and Candidate Arthritogenic Peptides Revealed through Heuristic Clustering of Data-independent Acquisition Mass Spectrometry (DIA-MS) Data. *Mol Cell Proteomics*. 2016;15(6):1867-76.
- Nossent JC, Johnsen S, Bakland G. The influence of ERAP1 gene variants on clinical phenotype in ankylosing spondylitis. *Scand J Rheumatol*. 2016;45(6):474-9.
- Su W, Du L, Liu S, Deng J, Cao Q, Yuan G, et al. ERAP1 /ERAP2 and RUNX3 polymorphisms are not associated with ankylosing spondylitis susceptibility in Chinese Han. *Clin Exp Immunol*.2018;193(1):95-102.
- Tang Y, Yang P, Wang F, Xu H, Zong SY. Association of polymorphisms in ERAP1 and risk of ankylosing spondylitis in a Chinese population. *Gene*.2018;646:8-11.
- Han R, Xia Q, Xu S, Fan D, Pan F. Interleukin-23 receptor polymorphism (rs10889677 A/C) in ankylosing spondylitis: Meta-analysis in Caucasian and Asian populations. *Clin Chim Acta*.2018;477:53-9.
- Zhong L, Wang W, Song H. Complex role of IL-23R polymorphisms on ankylosing spondylitis: a meta-analysis. *Expert Rev Clin Immunol*. 2018;14(7):635-43.
- Gravallese EM, Schett G. Effects of the IL-23-IL-17 pathway on bone in spondyloarthritis. *Nat Rev Rheumatol*. 2018;14(11):631-40.
- Chen S, Li Y, Deng C, Li J, Wen X, Wu Z, et al. The associations between PD-1, CTLA-4 gene polymorphisms and susceptibility to ankylosing spondylitis: a meta-analysis and systemic review. *Rheumatol Int*. 2016;36(1):33-44.
- Dahmani CA, Benzaoui A, Amroun H, Mecabih F, Sediki FZ, Zemani-Fodil F, et al. Association of the HLA-B27 antigen and the CTLA4 gene CT60/rs3087243 polymorphism with ankylosing spondylitis in Algerian population: A case-control study. *Int J Immunogenet*. 2018;45(3):109-17.
- Cetintepe SP, Senturk T, Sargin G, Aydin N. Serum sCTLA-4 levels and clinical manifestations in

- ankylosing spondylitis patients. *Eur J Rheumatol*. 2018;5(2):115-7.
19. Wu J, Zhang L, Zhou YX. The association between CTLA-4(+49 A/G) polymorphism and susceptibility to ankylosing spondylitis: a meta-analysis. *Int J Rheum Dis*. 2016;19(12):1237-43.
 20. Wang NG, Wang DC, Tan BY, Wang F, Yuan ZN. TNF-alpha and IL10 polymorphisms interaction increases the risk of ankylosing spondylitis in Chinese Han population. *Int J Clin Exp Pathol*. 2015;8(11):15204-9.
 21. Mordovskii V, Semenchukov A, Nikulina SY, Salmina AB, Chernova A, Kapustina E, et al. TNFR1 -383 A>C polymorphism and ankylosing spondylitis in a Russian Caucasian population: a preliminary study. *Genet Mol Res*. 2017;16(1).
 22. Hu N, Cui Y, Yang Q, Wang L, Yang X, Xu H. Association of polymorphisms in TNF and GRN genes with ankylosing spondylitis in a Chinese Han population. *Rheumatol Int*. 2018;38(3):481-7.
 23. Nossent JC, Sagen-Johnsen S, Bakland G. Tumor necrosis factor-alpha promoter -308/238 polymorphism association with less severe disease in ankylosing spondylitis is unrelated to serum TNF-alpha and does not predict TNF inhibitor response. *J Rheumatol*. 2014;41(8):1675-82.
 24. Sode J, Bank S, Vogel U, Andersen PS, Sorensen SB, Bojesen AB, et al. Genetically determined high activities of the TNF-alpha, IL23/IL17, and NFkB pathways were associated with increased risk of ankylosing spondylitis. *BMC Med Genet*. 2018;19(1):165.
 25. Xia Y, Liang Y, Guo S, Yu JG, Tang MS, Xu PH, et al. Association between cytokine gene polymorphisms and ankylosing spondylitis susceptibility: a systematic review and meta-analysis. *Postgrad Med J*. 2018;94(1115):508-16.
 26. Xu H, Qu Y. Correlation of PON1 polymorphisms with ankylosing spondylitis susceptibility: A case-control study in Chinese Han population. *Medicine (Baltimore)*. 2017;96(42):e7416.